## Viekira Pak® (ombitasvir, paritaprevir, ritonavir 12.5/75/50 mg; dasabuvir 250 mg) & Ribavirin 24 week Treatment Checklist

Prior to Treatment	
Labs	
Immediately prior: Pregnancy test (if applicable	)
Uric Acid	
Within 1 month: CBC	
CMP (If GFR <50, do not star	t treatment; consult Liver Disease Specialist)
PT/INR	
Within 3 months: HCV RNA	
Genotype confirmation	
Within 6 months: AFP	
TSH	
A1C or Fasting Glucose	
Vitamin D 250H (treat if def	icient)
Within 1 year: HIV screening	
Miscellaneous	
Hepatitis A status/screening if not done	
Hepatitis B status/screening if not done	
PHQ-9 baseline	
AUDIT-C	
Symptoms Inventory baseline	
Week 2 (with ribavirin)	
CBC	
CMP <sup>1</sup>	
Symptoms Inventory	3 months post treatment
,	CBC
Week 4	Liver Function Tests
HCV RNA	HCV RNA
CBC	PHQ-9
$\_\_$ CMP <sup>1</sup>	
Symptoms Inventory	
<pre> Pregnancy test (if applicable)</pre>	
	Nurse follow-up in clinic or by phone:
Weeks 8, 12, 16, & 20	Symptoms Inventory
CBC	Managing side effects
$\longrightarrow$ CMP <sup>1</sup>	Medication adherence discussion
Symptoms Inventory	Alcohol intake
<pre> Pregnancy test (if applicable)</pre>	Birth control reminder
Week 24	Refill reminder
HCV RNA	
CBC	
CMP <sup>1</sup>	
Symptoms Inventory	
<pre> Pregnancy test (if applicable)</pre>	

1- If GFR <50, consult Liver Disease Specialist.

Viekira Pak® (ombitasvir, paritaprevir, ritonavir 12.5/75/50 mg; dasabuvir 250mg) & Ribavirin 24 week Lab Tracking Form

General Patient Information	Pre-Treatment Lab Results	Medication Regimen					
Name:	HCV RNA:	1- Viekira Pak <sup>®</sup> Do not change dose. 2 pink tablets of ombitasvir, paritaprevir, ritonavir with breakfast.					
DOB:/	Genotype: HIV: TSH:	1 beige tablet of dasabuvir with breakfast and 1 with dinner.					
MRN:	Vit D 250H: AFP: GFR:	2- Ribavirin: mg/day PO divided into 2 doses. Take with breakfast & dinner. ≥75kg = 1200mg/day <75kg = 1000mg/day					
Phone #:	PT/INR: A1C/Glucose:	**Dose Reduction/Date:/					
Treatment Start Date:		**Additional Dose Change/Date:/**Consult ANTHC Liver Disease & Hepatitis Specialists for further guidance about dose changes.					

Completed														
Treatment Week	Lab Date	Hgb	Hct	WBC	PLT	ALT	AST	Alk Phos	Total Bili	Creat/ GFR	PHQ-9 (Specified weeks)	HCV RNA (Specified weeks)	Weight (kg)	Pregnancy Test
	Lub Dute	11,52	1100	1100		ALI	AJI	AIKTHOS	Total Bill	Creaty Grit	(Specified weeks)	(Specified weeks)	(1.6)	rest
Pre-Treatment														
Treatment Start Week 0											DUI 0 0	1101/ 5514		
vveek 0											PHQ-9	HCV RNA		
optional														
Week 2														
optional														
Week 4												HCV RNA		
optional														
optional														
Week 8														
optional														
optional														
Week 12											PHQ-9	HCV RNA		
optional														
Week 16														
optional														
Week 20														
optional														
Week 24											PHQ-9	HCV RNA		
3 months post														
treatment											PHQ-9	HCV RNA		1

Labs recommended for each follow up visit: CBC, CMP, pregnancy test (females of childbearing age), and HCV RNA as specified.

Please note the following critical values. These may require modification of dosage or discontinuation of causative med. Contact ANTHC Liver Disease Specialists with any questions.

Hgb <10.0 gm/dL If hemoglobin drops below 10, reduce ribavirin dose to 600mg (refer to ribavirin package insert). If hemoglobin <8.5, hold ribavirin & consult ANTHC Liver Disease Specialists.

GFR <50 If GFR is <50, decrease ribavirin dose (refer to ribavirin package insert) and consult ANTHC Liver Disease Specialists.

PLTs <50 K/uL If platelet count drops below 50, consult ANTHC Liver Disease Specialists.