Viekira Pak® (ombitasvir, paritaprevir, ritonavir 12.5/75/50 mg; dasabuvir 250 mg) 12 week Treatment Checklist

(with and without ribavirin)

Prior to Treatment		
Labs		
Immediately prior	: Pregnancy test (if applic	able)
	Uric Acid (with ribavirin)	
Within 1 month:	CBC with differential	
	CMP (If GFR <50, do not	start treatment; consult Liver Disease Specialist)
	PT/INR	
Within 3 months:	HCV RNA	
	Genotype confirmation	
Within 6 months:	AFP	
	TSH	
	A1C or Fasting Glucose	
	Vitamin D 250H (treat if	f deficient)
Within 1 year:	HIV screening	
Miscellaneous		
Hepatitis A	status/screening if not done	
	status/screening if not done	
PHQ-9 base		
AUDIT-C		
	Inventory baseline	
	,	
Week 2 (with ribavirin) CBC CMP¹ Symptoms Week 4 HCV RNA		3 months post treatment CBC Liver Function Tests HCV RNA PHQ-9
$___$ CBC $___$ CMP 1		Nurse follow-up in clinic or by phone:
	Inventory	Symptoms Inventory
Symptoms	test (if applicable)	Managing side effects
Fregulaticy	test (II applicable)	Medication adherence discussion
Week 8		Alcohol intake
CBC		Birth control reminder
CMP ¹		Refill reminder
Symptoms	Inventory	
	test (if applicable)	
11egnancy	test (ii applicable)	
Week 12		
HCV RNA		
CBC		
CMP ¹		
Symptoms	Inventory	
	test (if applicable)	
0 -7	, ,,	

1- If GFR <50, consult Liver Disease Specialist.

Viekira Pak® (ombitasvir, paritaprevir, ritonavir 12.5/75/50 mg; dasabuvir 250 mg) & Ribavirin 12 week Lab Tracking Form

Conoral	Dationt	Information
senerai	Patient	intormation

Pre-Treatment Lab Results

Medication Regimen

Name: DOB:/ MRN: Phone #:	HCV RNA: Genotype: HIV: TSH: Vit D 25OH: AFP: GFR:	1- Viekira Pak [®] Do not change dose. 2 pink tablets of ombitasvir, paritaprevir, ritonavir with breakfast. 1 beige tablet of dasabuvir with breakfast and 1 with dinner. 2- Ribavirin: mg/day PO divided into 2 doses. Take with breakfast & dinner. ≥75kg = 1200mg/day <75kg = 1000mg/day **Dose Reduction/Date: / **Additional Dose Change/Date: /				
Treatment Start Date:	PT/INR: A1C/Glucose:	**Consult ANTHC Liver Disease & Hepatitis Specialists for further guidance about dose changes.				

Completed											DUO 0			
Treatment								411 51			PHQ-9 (Specified	HCV RNA	Weight	Pregnancy
Week	Lab Date	Hgb	Hct	WBC	PLT	ALT	AST	Alk Phos	Total Bili	Creat/GFR	weeks)	(Specified weeks)	(kg)	Test
Pre-Treatment														
Treatment Start Week 0											PHQ-9	HCV RNA		
optional														
Week 2														
optional														
Week 4												HCV RNA		
optional														
optional														
Week 8														
optional														
optional														
Week 12											PHQ-9	HCV RNA		
optional														
3 months post treatment											PHQ-9	HCV RNA		

Labs recommended for each follow up visit: CBC, CMP, pregnancy test (females of childbearing age), and HCV RNA as specified.

Please note the following critical values. These may require modification of dosage or discontinuation of causative med. Contact ANTHC Liver Disease Specialists with any questions.

Hgb <10.0 gm/dL If hemoglobin drops below 10, reduce ribavirin dose to 600mg (refer to ribavirin package insert). If hemoglobin <8.5, hold ribavirin & consult ANTHC Liver Disease Specialists.

GFR <50 If GFR is <50, decrease ribavirin dose (refer to ribavirin package insert) and consult ANTHC Liver Disease Specialists.

PLTs <50 K/uL If platelet count drops below 50, consult ANTHC Liver Disease Specialists.